

Do attractive bosons condense?

N.K. Wilkin, J.M.F. Gunn and R.A. Smith

*School of Physics and Space Research,
University of Birmingham,
Edgbaston, Birmingham,
B15 2TT,
United Kingdom.
(February 1, 2008)*

Motivated by experiments on bosc atoms in traps which have attractive interactions (e.g. ^7Li), we consider two models which may be solved exactly. We construct the ground states subject to the constraint that the system is rotating with angular momentum proportional to the number of atoms. In a conventional system this would lead to quantised vortices; here, for attractive interactions, we find that the angular momentum is absorbed by the centre of mass motion. Moreover, the state is *uncondensed* and is an example of a ‘fragmented’ condensate discussed by Nozières and Saint James. The same models with *repulsive* interactions are fully condensed in the thermodynamic limit.

One of the most novel aspects of the creation of Bose condensates with neutral atoms in traps is the possibility of observing a bosc gas with *attractive* interactions (negative scattering lengths). The case of ^7Li has been studied both experimentally [1,2] and theoretically. Condensation has been predicted to be stable for a sufficiently small number of particles or sufficiently weak interactions [3,4]. The instability to collapse when these conditions are not obeyed has also been discussed by several authors [5–9].

In this Letter we show, using two exactly soluble models, that there may be other possibilities for non-condensed states with attractive interactions. The states are the ‘fragmented’ condensates discussed by Nozières and Saint James [10] in the context of excitonic bosc condensates. The possibility of such states emerges from the realisation [11] that it is the exchange interaction which causes bosons with *repulsive* interactions to condense into a single one-particle state, if there are several one-particle ground states. Conversely for attractive interactions, the exchange term is negative and may prefer ‘fragmented’ [10] condensation into more than one state if there is a degeneracy (or perhaps if the interactions are sufficiently strong). Kagan *et al.* [4] argue that trapped gases with sufficiently large negative scattering lengths are unstable to the formation of clusters using a somewhat different argument, but with the same physical origin.

The two models we examine are: particles in a harmonic trap with L quanta of angular momenta with attractive interactions treated as a degenerate perturbation [12]; rotating particles in a harmonic trap interacting with harmonic interactions [13–16]. (Both of these cases have been of interest for *fermions* [12,14], where rotation is replaced by a magnetic field and the phenomena are related to the fractional quantum Hall effect.) Rotation is considered in both cases, partly because the non-rotating ground state, in the thermodynamic limit, is trivial in both cases (for different reasons) and partly

because the response to rotation is characteristic of superfluidity in the system [17,18].

Consider the two-dimensional Hamiltonian

$$\mathcal{H} = \mathcal{H}_0 + \mathcal{H}_1$$

$$\mathcal{H}_0 = -\frac{1}{2}\nabla^2 + \frac{1}{2}\sum_i \mathbf{r}_i^2 \text{ and } \mathcal{H}_1 = \frac{\eta}{2}\sum_{i>j} \delta(\mathbf{r}_i - \mathbf{r}_j), \quad (1)$$

in the limit where the dimensionless coupling is weak, $|\eta| \ll 1$, so that the contact interaction can be treated perturbatively. We will now determine the ground state subject to the constraint that the system contains L quanta of angular momentum. We should note that the centre of mass variables will separate in this Hamiltonian because the trap is harmonic. This will be used below.

The single particle spectrum is usefully expressed [19] in terms of the angular momentum quantum number, m , and the radial quantum number, n_r :

$$E = |m| + 2n_r + 1 \quad (2)$$

in dimensionless units. For the non-interacting case, $\eta = 0$, with N particles it is clear that to minimise the energy $n_r = 0$ for all particles. The angular momentum may be expressed as $L = \sum_m mN_m$ where N_m is the number of particles in the state m . There is a degeneracy, in general, associated with the choice of the set $\{N_m\}$. If the angular momentum is positive, $L > 0$, then because of the modulus signs in Eq. (2) the energy is minimised if one takes only *positive* integers for m . Then for $0 < |\eta| \ll 1$ degenerate perturbation theory should be used with a basis set consisting of all states $\{N_m\}$, such that $L = \sum_{m=0}^L mN_m$.

The single particle states with $n_r = 0$ and $m \geq 0$ are of the form $z^m \exp(-|z|^2/2)$ where $z = x + iy$ and m is the angular momentum quantum number. With the use of a computer algebra package we have been able to study systems of up to six particles comprehensively. In

addition we can prove that the form of the ground state holds for an arbitrary number of particles.

We find that in all cases the groundstate for the attractive interaction ($\eta < 0$) is $\psi_{z_c} = z_c^L \exp \sum_{i=1}^N (-|z_i|^2)$ where $z_c = \sum_{i=1}^N z_i$, is the centre of mass and L is the total angular momentum in the system. The contact interaction energy contribution for the groundstate is independent of L , $\epsilon(N, L) \propto \eta N(N-1)/2$. To prove the form of the groundstate wavefunction we show that ψ_{z_c} is the unique eigenfunction of $V = \sum_{i < j} \delta(\mathbf{r}_i - \mathbf{r}_j)$ corresponding to its largest eigenvalue, $\lambda_{\max}$. First we note that ψ_{z_c} is trivially an eigenfunction of V when $L = 0$ since it is the only state in the $L = 0$ subspace. Since the centre of mass coordinate, z_c , can be separated out in the Hamiltonian, it follows that ψ_{z_c} is an eigenfunction for any L .

Let us now work in the basis

$$|m_1, m_2 \dots m_N\rangle = \prod_{i=1}^N z_i^{m_i} e^{-|z_i|^2/2}, \quad (3)$$

where $\sum m_i = L$. The matrix elements $\langle m|V|m'\rangle$ are non-negative, and are positive when $m_i + m_j = m'_k + m'_l$ for some i, j, k, l , and $m_p = m'_q$ for the remaining labels. The coefficients $\langle m|\psi_{z_c}\rangle$ are all positive, from which it follows that the eigenvector ψ_{z_c} belongs to the largest eigenvalue $\lambda_{\max}$. To see this note that $\lambda_{\max}$ can be derived from variational principle

$$\lambda_{\max} = \max \left\{ \frac{\langle \psi|V|\psi \rangle}{\langle \psi|\psi \rangle} \right\} = \max \left\{ \frac{\sum_{ij} V_{ij} \psi_i \psi_j}{\sum_i \psi_i^2} \right\}. \quad (4)$$

If we take an eigenvector of $\lambda_{\max}$, and replace all its components by their absolute values, the variational functional in Eq. (4) cannot decrease, and so must remain at $\lambda_{\max}$. It follows that there is an eigenvector of $\lambda_{\max}$ whose components are non-negative; ψ_{z_c} has non-zero overlap with this eigenvector, and so must belong to $\lambda_{\max}$.

To prove non-degeneracy of the eigenspace of $\lambda_{\max}$ we note that the matrix V_{ij} is ‘‘connected’’ in the following way: if we take a basis vector $|i\rangle$ and consider all $|j\rangle$ with $V_{ij} > 0$, then consider all $|k\rangle$ with $V_{jk} > 0$ and so on, this includes all basis vectors. If the eigenspace of $\lambda_{\max}$ is degenerate then there must be an eigenvector of $\lambda_{\max}$ whose components are of both signs (since we can choose this eigenvector to be orthogonal to ψ_{z_c}), and all non-zero (if some components are zero, simply add on a very small amount of ψ_{z_c}). The vector made by taking the absolute value of the latter’s components will also be an eigenvector. The difference in value of the variational functional for the two vectors can only be zero if the i -th and j -th components have same sign when $V_{ij} > 0$. By connectedness we see that this means all components must have the same sign, and hence the eigenspace is non-degenerate. This completes the proof that ψ_{z_c} is the non-degenerate ground state of the Hamiltonian Eq. (1) with attractive interaction.

To determine the degree of condensation, if any, the single particle density matrix, $\rho(z, z'^*)$ is required for the ground state. Yang [20] showed that off-diagonal long-range order is associated with the largest eigenvalue of the density matrix (the magnitude of the eigenvalue is the fraction condensed) with the ‘condensate wavefunction’ being the associated eigenvector. The notion of off-diagonal long-range order is not of such great use for trapped atoms, but this definition of the *condensate* is useful in an inhomogeneous setting. The single particle density matrix has the form:

$$\rho(z, z'^*) = \frac{1}{Q} \int \prod_{i=2}^N dz_i dz_i^* \psi(z, z_2, \dots z_N) \psi^*(z', z_2, \dots z_N), \quad (5)$$

where Q is the normalisation. On integrating we find,

$$\rho(z, z'^*) = \frac{e^{-|z|^2/2} e^{-|z'|^2/2}}{\pi} \sum_{m=0}^L z^m z'^{*m} \frac{(N-1)^{L-m} L!}{N^L (L-m)! m!^2}. \quad (6)$$

Thus the resulting eigenfunctions and eigenvalues for a given m are

$$\psi_m = e^{-\frac{|z|^2}{2}} z^{*m} \quad \text{and} \quad \rho_m = \frac{(N-1)^{L-m} L!}{N^L (L-m)! m!}. \quad (7)$$

If we now consider the case of $L = Nq$ (which in a conventional system, *e.g.* ^4He would correspond to q vortices) then if a condensate exists its eigenvalue will correspond to $m = q$. Simplifying we find

$$\rho_q = (1 - 1/N)^{(qN-q)} \frac{(Nq)!}{N^q (qN-q)! q!}, \quad (8)$$

which can be rewritten as a Poisson distribution in the limit that $N \rightarrow \infty$. On taking the further limit of $q \rightarrow \infty$ the maximal eigenvalue becomes $\rho_q \sim 1/\sqrt{2\pi q}$. However, the eigenvalues of significant weight are distributed over $q - \sqrt{q} \lesssim m \lesssim q + \sqrt{q}$. This is clearly not the pronounced peak required for a condensate and is reminiscent of Nozières and Saint James’ fragmented condensate [10]. Lest this be thought to be misleading for small q , we note the following results for $q = 1$. We find that the eigenvalue where the putative ‘condensate’ would be, $\rho_1(q = 1) = e^{-1}$, that $\rho_0(q = 1) = e^{-1}$ as well and that $\rho_2(q = 1) = \frac{1}{2}e^{-1}$. The ‘condensate’ is not singled out as having a uniquely large eigenvalue.

The first excited state, Φ , is also of interest, as we find a rudimentary ‘vortex’. The general form is:

$$\Phi = z_c^{L-2} \sum_{i>j}^N (z_i - z_j)^2 \quad \text{with} \quad \epsilon_1 \propto N(N-2)/2. \quad (9)$$

Again from symmetry considerations we find that we require a minimum of 2 quanta of angular momentum in

order to produce an excited state. For $L=2$, we find that there are two possible states for all N and these correspond to the groundstate and excited state we have described above. Because of the separation of the centre of mass variables mentioned above, this is in fact a general result: for $\psi_{z_c} = z_c^L$ then an excited state is Φ (although we have not proved that it is always the *first* excited state).

To determine whether the results from the contact interaction model are likely to be generic or are artifacts (for instance of degenerate perturbation theory), we turn to the second model. The Hamiltonian [15,16] (first discussed in the context of nuclear physics [13]) describes N bosons with attractive harmonic coupling ($\tilde{\Lambda} > 0$) (i labels the particles):

$$H = -\frac{\hbar^2}{2m} \sum_i^N \tilde{\nabla}_i^2 + \frac{k}{2} \sum_i^N \mathbf{x}_i^2 + \frac{\tilde{\Lambda}}{4} \sum_{i,j}^N (\mathbf{x}_i - \mathbf{x}_j)^2, \quad (10)$$

where we enforce the symmetry of the wavefunctions at the end of the calculation. In dimensionless units, $\Lambda = \tilde{\Lambda}/k$, $y = (\hbar^2/(mk))^{1/4} x$ and $\nabla = (mk/\hbar^2)^{1/4} \tilde{\nabla}$

$$\mathcal{H} = -\frac{1}{2} \sum_i^N \nabla_i^2 + \frac{1}{2} \sum_i^N \mathbf{y}_i^2 + \frac{\Lambda}{4} \sum_{i,j}^N (\mathbf{y}_i - \mathbf{y}_j)^2, \quad (11)$$

which upon rearrangement leads to

$$\mathcal{H} = -\frac{1}{2} \sum_i^N \nabla_i^2 + \frac{(1 + N\Lambda)}{2} \sum_i^N \mathbf{y}_i^2 - \frac{\Lambda}{2} \left(\sum_i^N \mathbf{y}_i \right)^2. \quad (12)$$

Here we note that the problem in d dimensions separates into d one-dimensional problems. Hence we will now restrict ourselves to one-dimension for clarity.

To determine the degree of condensation we again need to calculate the single particle density matrix. To do this we change variables to the centre of mass coordinate, $\zeta_N = 1/\sqrt{N} \sum_i^N y_i$, and ζ_i , ($i = 1, \dots, N-1$) which are arbitrary but orthogonal to ζ_N . The Hamiltonian in these variables is

$$\mathcal{H} = -\frac{1}{2} \nabla_{\zeta}^2 + \frac{(1 + N\Lambda)}{2} \sum_i^N \zeta_i^2 - \frac{N\Lambda}{2} \zeta_N^2, \quad (13)$$

This leads to the groundstate wavefunction having the form

$$\psi = e^{-\frac{1}{2}(1+N\Lambda)^{1/2} \zeta^2} e^{-\frac{1}{2}(1-(1+N\Lambda)^{1/2}) \zeta_N^2}, \quad (14)$$

where $\zeta = \{\zeta_1, \dots, \zeta_N\}$. The corresponding frequencies are $\epsilon_i = (1 + N\Lambda)^{1/2}/2$ for $i \neq N$ and $\epsilon_N = 1/2$. We now consider the model in $d \geq 2$ with attractive interactions in the fixed angular momentum subspaces. In the groundstate we find that all angular momentum is absorbed by the centre of mass variable. This may be seen

by noting that the centre of mass oscillators (associated with the different components of the centre of mass motion) have a lower associated frequency than the other oscillators describing relative motion. The physical interpretation is straightforward: relative motion requires more energy as work must be performed against the attractive interactions. (The converse will hold true for repulsive interactions). Hence, for two-dimensions and L quanta of angular momentum we can immediately write down the wavefunction,

$$\psi = z_c^L e^{-\frac{1}{2}(1+N\Lambda)^{1/2} |\mathbf{z}|^2} e^{-\frac{1}{2}(1-(1+N\Lambda)^{1/2}) |z_c|^2}. \quad (15)$$

In the thermodynamic limit it can be shown that the contribution to the single particle density matrix of the exponential term associated with z_c is negligible. Hence, surprisingly, the density matrix reduces to that of the contact interaction model. The groundstate wavefunctions will therefore be the same, as will the properties of the single particle density matrix.

We shall now show that these systems are condensed (at least under some conditions) when the interactions are repulsive. Thus the lack of condensation is not due to peculiarities of the models in general, but of the attractive interactions in particular.

The repulsive case has also been considered, and will be discussed elsewhere. We find no vortex lattice, although a Laughlin like state exists for some special values of L .

Firstly consider the contact interaction model when there are N quanta of angular momentum in the system. Conventionally there would then be one vortex in the system, and one might expect that the ground state (subject to the constraint of the amount of angular momentum) would be:

$$\psi_{L=N}^{\text{mft}} = \prod_{i=1}^N \left(z_i e^{-|z_i|^2/2} \right) \quad (16)$$

We conjecture the following form for the CIM:

$$\psi_{L=N}^{\text{exact}} = \prod_{i=1}^N \left([z_1 + \dots + z_n - N z_i] e^{-|z_i|^2/2} \right) \quad (17)$$

We have demonstrated that this form is correct by explicit calculation on systems of up to 6 bosons. The physical interpretation of this wavefunction is that the bosons are rotating around the centre of mass, which would be a condensate if the centre of mass were a c -number. We will now show that in the thermodynamic limit the corrections to full condensation are $O(1/N)$. Consider the density matrix constructed from the wavefunction Eq. (17):

$$\rho(z, z'^*) = e^{-|z|^2/2} e^{-|z'|^2/2} \int \left\{ \prod_{i=2}^N dz_i dz_i^* \right\} \times \left(\omega - \frac{(N-1)}{N} z \right) \left(\omega^* - \frac{(N-1)}{N} z'^* \right) \times \prod_{j=2}^N \left\{ \left(\omega + \frac{z}{N} - z_j \right) \left(\omega^* + \frac{z'^*}{N} - z_j^* \right) e^{-|z_j|^2} \right\} \quad (18)$$

where $\omega = 1/N \sum_{i=2}^N z_i$. Then to separate the integration over the different z_i we introduce a delta function for the centre of mass variable:

$$1 = \int_{-\infty}^{\infty} d\omega_x d\omega_y \delta \left(\omega_x - \frac{1}{N} \sum_{i=2}^N x_i \right) \delta \left(\omega_y - \frac{1}{N} \sum_{i=2}^N y_i \right) \quad (19)$$

and use the integral representation

$$\delta \left(\omega - \frac{1}{N} \sum_{i=2}^N \mathbf{r}_i \right) = \frac{1}{(2\pi)^2} \int d\boldsymbol{\lambda} e^{i\boldsymbol{\lambda} \cdot \left(\omega - \frac{1}{N} \sum_{i=2}^N \mathbf{r}_i \right)} \quad (20)$$

Upon substitution and integration we find that, in the limit that $N \rightarrow \infty$, to accuracy $O(1/N^2)$:

$$\rho(z, z'^*) = \sum_n \psi(z) \rho_n \psi^*(z') = e^{-|z|^2/2} \left(\frac{1}{N} \frac{1}{\sqrt{\pi}} \frac{1}{\sqrt{\pi}} + \left[1 - \frac{2}{N} \right] \frac{z}{\sqrt{\pi}} \frac{z'^*}{\sqrt{\pi}} + \frac{1}{N} \frac{z^2}{\sqrt{2\pi}} \frac{z'^{*2}}{\sqrt{2\pi}} \right) e^{-|z'|^2/2} \quad (21)$$

Thus there is a condensate with eigenvalue $1 - 2/N$, in the state z , which is fully condensed in the thermodynamic limit. The corrections of $1/N$ are in the states 1 and z^2 . In addition a Laughlin state is empirically found in small systems to be the ground state for $L = N(N-1)$.

Turning to the harmonic interactions model, we note that the centre of mass oscillator has a higher frequency than the others when the interaction is repulsive. In that case the other oscillators will be populated in preference when minimising the energy subject to L (a multiple of N) quanta of angular momentum. Now, the other oscillators are degenerate and the centre of mass factor is not being multiplied by a large multiple of the centre of mass coordinate, which it was in the attractive case. The latter implies that the centre of mass factor is irrelevant in the thermodynamic limit. Thus the ground state reduces to a single particle form [Eq. (2)] and hence the answer will be a condensate into the state $z^{L/N}$.

In conclusion we have shown that for a model with degenerate ground states in the absence of interaction, there is no condensate formed when weak interactions are incorporated. Consequently, in this particular case, there is no vortex lattice. In a different model, which

does not have degenerate ground states, we have shown that the particles are uncondensed when given an extensive quantity of angular momentum. In both cases the angular momentum of the system resides in the centre of mass motion, in contrast to the more familiar case of repulsive interactions. This leads to the general hypothesis: attractive bosons do not condense in the presence of single particle degeneracy and their angular momentum resides in the centre of mass motion. The investigation of *rotating* ^7Li might be fruitful in exposing an uncondensed 'ground state'. Repulsive interactions in the same models lead to condensed ground states, so showing that attractive interactions do indeed lead to different physics.

We would like to thank M.W. Long, D.A. Lowe and A.G.B. Triulzi for useful discussions and EPSRC for financial support through grants GR/J35238, GR/L28784 and GR/L29156.

-
- [1] C. Bradley, C. Sackett, J. Tollett, and R. Hulet, Phys. Rev. Lett. **75**, 1687 (1995).
 - [2] C. Bradley, C. Sackett, and R. Hulet, Phys. Rev. Lett. **78**, 985 (1997).
 - [3] P. Ruprecht, M. Edwards, K. Burnett, and C. Clark, Phys. Rev. A **54**, 4178 (1996).
 - [4] Y. Kagan, G. Shlyapnikov, and J. Walraven, Phys. Rev. Lett. **76**, 2670 (1996).
 - [5] G. Baym and C. Pethick, Phys. Rev. Lett. **76**, 6 (1996).
 - [6] R. Dodd, M. Edwards, C. Williams, C. Clark, M. Holland, P. Ruprecht, and K. Burnett, Phys. Rev. A **54**, 661 (1996).
 - [7] E. Shuryak, Phys. Rev. A **54**, 3151 (1996).
 - [8] L. Pitaevskii, Phys. Lett. A **221**, 14 (1996).
 - [9] V. Babichenko and A. Kozlov, JETP Lett. **64**, 729 (1996).
 - [10] P. Nozières and D. Saint James, J. Phys. (Paris) **43**, 1133 (1982).
 - [11] P. Nozières, in *Bose-Einstein Condensation*, edited by A. Griffin, D. W. Snoke, and S. Stringari (CUP, Cambridge, 1995), Chap. 2.
 - [12] S. Trugman and S. Kivelson, Phys. Rev. B **31**, 5280 (1985).
 - [13] S. Gartenhaus and C. Schwartz, Phys. Rev. **108**, 482 (1957).
 - [14] S. Girvin and T. Jach, Phys. Rev. B **28**, 4506 (1983).
 - [15] N. Johnson and M. Payne, Phys. Rev. Lett. **67**, 1157 (1991).
 - [16] F. Brosens, J. T. Devreese, and L. F. Lemmens, preprint cond-mat/9704040.
 - [17] A. J. Leggett, Physica Fennica **8**, 125 (1973).
 - [18] P. Nozières and D. Pines, *The Theory of Quantum Fluids* (Addison-Wesley, Redwood City, California, 1990), Vol. II.
 - [19] S. Flugge, *Practical Quantum Mechanics* (Springer Verlag, Berlin, 1974), p109.
 - [20] C. N. Yang, Rev. Mod. Phys. **34**, 694 (1962).